

**Adopted Regulation Strategy
Lake of the Woods Control Board Secretariat
March 10, 2015**

The Lake of the Woods Control Board held a Regulation Meeting on March 10, 2015 in Kenora and adopted a regulations strategy for the spring. The strategy was formulated considering basin conditions, hydrological and meteorological forecasts, and the input of the various interests concerned with basin management. Input was provided in written and verbal reports as well as from the Board's Regulation Guide (<http://www.lwcb.ca/reg-guide/index.html>).

For an update on current conditions, please refer to the Basin Data section of the Board's web site at <http://www.lwcb.ca/waterflowdata.html>. For regulation actions and directives taken under the strategy please see the Regulation Actions at <http://www.lwcb.ca/regulation/index.html>.

Lac Seul

A) Seasonal Considerations

The end-of-winter (April 15) level target range set at the October Regulation Meeting (354.7 m / 1163.7 ft) remains an achievable target under current conditions, and will likely require adjustments to outflow if freshet is early. This target level is approximately 4 cm more than the maximum winter drawdown of 1.5 m / 4.9 ft preferred for fisheries interests on Lac Seul.

To support spawning conditions and navigation in early spring, it is desirable to have the lake level rising after mid-April. This will likely require outflow reductions, but rising lake levels must be balanced against a future risk of high water. The points below reflect a number of ideal or desirable regulation objectives over the next few months, based on input provided to the Board.

- Regulation of Lac Seul level and outflow should take into account the preferred Lac Seul, Pakwash Lake and English River levels for the fishery and tourist outfitter interests, to provide good spring spawning conditions and adequate navigation levels at the start of the walleye fishing season.
- Lac Seul level should be constant or rising after April 15 for spring spawning fish.
- The minimum spring lake level should be no more than 1.5 m (5 ft) below the November 1 level for whitefish. For spring 2015, this level is 354.74 m (1163.85 ft).
- The desirable lake level on May 15 should be no less than 355.1 m (1165 ft) for the start of the walleye fishing season.
- The tourist outfitters preferred summer maximum level is 356.6 m (1170 ft).
- Lac Seul level and outflow should be managed to supply water requested by Ontario Power Generation and Manitoba Hydro for hydroelectric energy generation, to avoid spill in wet conditions and violation of low flow constraints in dry conditions.
- A minimum flow of 180 m³/s below Manitou Falls is desirable during spring spawning. If there is not sufficient water to meet this criteria, Lac Seul discharge should be set to maintain a uniform flow through the spawning period.
- River flows at Grassy Narrows should be less than 550 m³/s.
- Lac Seul storage should be used to offset Lake of the Woods high/low outflow for the benefit of users of the Winnipeg River in Manitoba.

- Lac Seul level and outflow should be managed to reduce the need to close the Lake St. Joseph diversion with resulting spill down the Albany River. However, the diversion should nevertheless be closed to reduce impacts in the English and Winnipeg River basins under wet conditions.
- If inflow is above normal by May 15 and Lac Seul level meets the preferred level on that date, adjust outflow in May and June so as to target a June 30 elevation of 356.3 m / 1169.0 ft, thereby reducing the risk of higher outflows in summer.

B) Strategy

To April 15 (Drawdown Period)

Incremental Lac Seul outflow adjustments should be made through March and early April in order to target an end-of-winter (April 15th) level of 354.7 m / 1163.7 ft. OPG has requested that the target level be reached as early as March 31 to accommodate an annual outage. This should be possible to achieve at the current drawdown rate and within the time remaining, provided that freshet is not too early. Outflow would then need to be adjusted to match inflow to hold the lake level relatively constant, until freshet begins.

After April 15 (Refill Period)

i) Low Inflow Conditions

- Outflow should be managed to ensure that Lac Seul level does not decline, and preferably rises, while providing sufficient outflow to meet downstream hydropower generation and fishery requirements.
- Target for an end-of-June level no lower than 355.46 m (1165.2 ft) with an outflow no lower than 100 m³/s, provided that flow in Manitoba is no lower than 600 m³/s.
- Target for an end-of-June level no lower than 355.1 m (1165.0 ft) with an outflow no lower than 25 m³/s, provided that flow in Manitoba is no lower than 300 m³/s.
- Consultation with interests, including OMNRF staff, tourist outfitters and the provincial hydro utilities, may be necessary to arrive at the appropriate balance between lake level and outflow.
- If inflow remains low throughout the refill period, outflow should be adjusted to maintain a balance between upstream and downstream interests. Note that a lower decile outflow for May for the 1981-2010 period was approximately 45 m³/s, while a 5th percentile outflow was below 35 m³/s.

ii) Moderate Inflow Conditions

- Due to concerns of high water, regulate through the refill period to hedge against wetter conditions, so that the risk of outflow above 500 m³/s is reduced.
- Generally target for lake levels between lower and upper quartile, while supplying water for hydropower production and for English River fishery concerns.
- Use additional water to maintain desired fishery flows in the English River below Manitou Falls, provided this does not cause high flow conditions on the Winnipeg River in Manitoba.
- Target for flow in the Winnipeg River in Manitoba between 675 and 960 m³/s.

iii) High Inflow Conditions

- As above, regulate through the refill period to hedge against a return to wetter conditions, so that the risk of outflow above 500 m³/s is reduced.
- Balance Ear Falls outflow with the rise in Lac Seul level to reduce flood risk both on Lac Seul and on downstream areas such as Pakwash Lake.
- Seek to maintain Lac Seul level (or projected level) below upper quartile through the refill period. Outflow should remain below 450 m³/s for moderately wet conditions, below 500 m³/s for most conditions, and below 600 m³/s in all but extreme conditions.
- Increase outflow to as much as 800 m³/s to keep the level below 357.1 m (1171.6 ft).
- When Lac Seul is above the level at which the Lake St. Joseph diversion comes under Board jurisdiction (356.01 m / 1168.0 ft until the end of May; 356.31 m / 1169.0 ft for June), the diversion flow should be reduced before increasing Lac Seul outflow above 500 m³/s.

Lake of the Woods

A) Seasonal Considerations

Lake of the Woods may not reach the target end-of-winter (March 31st) level of 322.35 m (1057.6 ft) set at the October Regulation Meeting due to the potential for an early freshet. The snowpack is beginning to melt but there remains considerable ice on the lake. The implementation of the proposed strategy for the upcoming period will look to the lessons learned during past freshet periods with similar conditions in an effort to provide effective regulation. Additionally, the risk of high water must be balanced against the risk of not meeting refill goals if spring rainfall is below normal.

The points below reflect a number of ideal or desirable regulation objectives over the next few months, based on input provided to the Board.

- Adjust lake level and outflow to achieve a balance between upstream and downstream interests, as inflow dictates.
- Minimize ice damage when possible. Ice damage is greater in the spring if there are rapid changes in water level (on either the lake or the river) and especially if the level rises while there is still a solid ice cover.
- The preferable end-of-April level for Lake of the Woods fishery is no lower than 322.5 m (1058.0 ft). Higher levels would be beneficial to northern pike.
- Regulate to avoid, to the extent possible, any reductions in outflow or any large increases in outflow during the spring spawning season on the Winnipeg River (late April to early June).
- For loons on the Winnipeg River, flow changes during the incubation period (approximately mid-May to the end of June) should be minimized.
- A summer Lake of the Woods level 10-15 cm (4-6 in) below the summer peak median level of 323.14 m (1060.2 ft) is desired for south shore residents. Satisfying these criteria would result in a summer peak level of about 323.0 m (1059.7 ft).
- For wild rice on Lake of the Woods and the Winnipeg River, maintain lower lake and river levels and minimize level and flow increases during the floating leaf stage in June and early July.
- For piping plovers on Lake of the Woods, maintain lower lake levels and minimize lake level increases during their nesting and rearing season of May, June and July.

- Within the regulation parameters for Lake of the Woods, regulate outflow to assist in meeting targets/preferences for the Winnipeg River in Manitoba.
- At the March Regulation Meeting, a number of interests on Lake of the Woods stated a preference for slightly lower lake levels in summer 2015 to assist with repairs to shorelines and structures that were damaged in 2014.

B) Strategy

Until March 31st (Drawdown Period)

Manage outflow to reach a March 31st level between lower quartile and median (322.38 to 322.49 m / 1057.7 to 1058.0 ft). Outflow should not be increased unless prolonged high inflow conditions are projected. Although the strategy set in October 2014 sought to target the lower end of this range, the signs indicating an early freshet suggest that the level might be closer to the upper end of this range. Should the freshet response be high enough in March to cause the lake to begin to rise appreciably, regulation will seek to balance lake level rise with outflow increases to attempt to limit the potential for ice damage to shoreline structures on both the lake and the river.

After March 31st (Refill Period)

Historically, the refill rate of Lake of the Woods is a factor more of the timing and magnitude of spring rainfall than of snowpack at the end of winter. Given the current conditions, the Board will strive to set outflows towards the high-normal range sooner when inflow is moderate or above so that river flows might not need to reach high rates for long periods. This would also position outflow to be increased or decreased relatively quickly in response to rainfall. Since spring rainfall is highly variable and precipitation forecasts are unreliable beyond horizons of more than a few days, regulation must be continually updated and outflow adjusted in response to changing conditions and forecasts.

i) Low Inflow Conditions

- Adjust outflow as necessary (subject to minimum flow requirements as recommended by the Ontario Ministry of the Environment) to keep the lake from declining. If possible, have the lake maintain at least a 10th percentile level, with outflow no lower than 100 m³/s.
- Assess conditions immediately before spawning begins in the Winnipeg River so that outflow can be set to prevent, as much as possible, the need for further flow reductions during the spawning season (late April to early June), while ensuring the lake level does not decline.
- Target a lake level above 322.78 m / 1059.0 ft at the end of June, if possible, with outflow no lower than 200 m³/s. This level corresponds to the lower end of the preferred range stated by the Lake of the Woods District Property Owners Association.
- If inflow remains low throughout the refill period, outflow should be adjusted to maintain a balance between upstream and downstream interests.

ii) Moderate Inflow Conditions

- Assess conditions immediately before spawning, as described under “Low Inflow Conditions” above.

- Outflow increases should be kept moderate during the spawning period and reductions should be minimized.
- Set outflow as high as 800 m³/s to prevent the peak lake level from exceeding 323.09 m / 1060 ft for the benefit of Lake of the Woods cottagers, if inflow is no higher than median.
- In accordance with the commitment made by the Board following the high water year of 2001, the lake level targets are 322.7 m (1058.7 ft) at the end of May and 322.9 m (1059.4 ft) at the end of June. Balance this by attempting to avoid outflow above the generation capability at Kenora and by optimizing hydroelectric generation downstream.
- Through late May and June, attempt to limit Lake of the Woods outflow changes that would adversely affect nesting loons on the Winnipeg River.
- Through June (and early July), try to manage lake levels to limit the rate of rise for wild rice.
- Maintain Nutimik Lake levels in the preferred range, to the extent possible, by managing outflow from both Lake of the Woods and Lac Seul.

iii) High Inflow Conditions

- Balance higher water levels on the lake with the impact of increased outflow downstream, both in Ontario and Manitoba.
- Avoid outflow above 800 to 900 m³/s if the lake level (or projected level) will remain below 323.2 m / 1060.4 ft in June. A flow of 900 m³/s on the Winnipeg River would cause the level below the Norman Dam to rise 1.4 m (4.6 ft) above normal, whereas a lake level of 323.2 m is less than 10 cm (4 in) above median by the end of June.
- Increase outflow as necessary to prevent the lake level (or the projected level) from rising above 323.47 m (1061.25 ft), which is the legislated top of the normal operating range
- An attempt should be made to keep outflow increases to a maximum of 100 m³/s per week, except during the spawning season when it would be desirable to not exceed 50 m³/s per week. Persistent higher inflow could, however, necessitate inflow increases of 200 m³/s per week or more.